

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

T G E 2 5 0 " X " P 3 2 5 0

Fig. 1.



α -glucosidase constructs

Fig. 2.A




Fig. 2.B.





Fig. 2.C.



Transcription Initiation site.

α s1 casein sequence, promoter or 3' untranslated region.

2 3 The boxes represent the exons in the α -glucosidase sequence, the thin line represents the intron sequences.

The numbers above the boxes are the exon numbers.

PA = polyadenylation signal.

ATG = translation initiation site.

TAG = translation stop codon

Fig. 3A.






Fig. 3B.



102250 " C 4 32260

Fig. 3.C.



□ = exon α-glu — = intron α-glu — = pKUN vector sequence

102250 16438860

Fig. 4. A.






Fig. 4. B.



202000 190000 180000 170000

Fig. 5.



20250 " 20260

Fig. 6.





Fig. 7.



T02250 "Z24h92260

Fig. 8.




Fig. 9.




Fig. 10.




Fig. 11. A.




Fig. 11. B.





Fig. 11. C.



Transgenic and non-transgenic whey on a Butyl column

Fig. 11. D.

0.998664427 0.9922027




Fig. 12.




Fig. 13. A.
transgenic whey




Fig. 13. B.
non-transgenic whey





Fig. 14.




12099801:1_UV1_280nm

12099801:1_pH

12099801:1_Cond%

12099801:1_Fractions

Fig. 15.



— 12099802:11_UV1_280nm

- - - 12099802:11_pH

* * * 12099802:11_Cond%

12099802:11_Fractions

Fig. 16.





Fig. 17.




12099804:13_UV1_280nm

12099804:13_pH

12099804:13_Cond%

12099804:13_Fractions

Fig. 18.




— 121099805:1_UV1_280nm

- - - 121099805:1_pH

* * * 121099805:1_Cond%

121099805:1_Fractions

Fig. 19.



— 121099806:1_UV1_280nm

- - - 121099806:1_pH

* * * 121099806:1_Cond%

121099806:1_Fractions

Fig. 20.




Fig. 21.

DATA SHEET - 062203





Fig. 22.

Chromatogram of Hatypei03:12



——— hatypei03:12_UV1_280nm

- - - - - hatypei03:12_pH

--* hatypei03:12_Cond%

hatypei03:12_Fractions

Fig. 23.

DEPARTMENT OF CHEMISTRY




Fig. 24.




Fig. 25.




Fig. 26.

XK16/15 80°C
cHT type I 10mM Napi pH 6.5 ; QFF eluate
Run 02249901/02259901/02269901

